arXiv:1204.1708v1 [quant-ph] 8 Apr 2012

Dynamics of coupled cavity arrays embedded in a non-Markovian bath

Xinyu Zhao[1]*, Jun Jing[1] J. Q. You[2], and Ting Yu[1]†
[1] *Center for Controlled Quantum Systems and Department of Physics and Engineering Physics, Stevens Institute of Technology, Hoboken, New Jersey 07030, USA and*
[2] *Department of Physics and State Key Laboratory of Surface Physics, Fudan University, Shanghai 200433, China*

In this paper, the non-Markovian dynamics of a coupled N-cavity model is studied based on the quantum state diffusion (QSD) approach. The time-local Diósi-Gisin-Strunz equation and the corresponding master equation are derived for a coupled cavity array. Non-Markovian effects are studied in two examples, two-cavity and three-cavity, under different boundary conditions. We have shown that the environment-memory can facilitate the cat-like state transfer from one cavity to another in the case of a strongly non-Markovian environment. We show that the non-Markovian QSD approach can be a valuable tool for the dynamics of a multi-partite continuous-variable (CV) system.

PACS numbers: 42.50.Lc, 03.65.Yz, 05.30.-d

I. INTRODUCTION

The dynamics of quantum open systems is one of the most important research topics in quantum optics, quantum dissipative system and quantum information [1, 2]. Open system dynamics is also essential for a deeper understanding of decoherence and implementation of quantum control [3–12]. While the theory of open systems in the Markov regime is well understood using the standard Lindblad master equations or the corresponding Markov quantum trajectories [13–16], the recent surge of interest in non-Markovian phenomena requires a set of systematic theoretical tools for approaching quantum dynamics in strong non-Markovian regimes. Among many recent research efforts in developing a theoretical approach to investigating the non-Markovian evolution, the non-Markovian quantum state diffusion (QSD) equation initially proposed by Diósi, Gisin, and Strunz [17, 18] provides a very powerful tool to deal with quantum open systems coupled to a bosonic environment. The Diósi-Gisin-Strunz QSD equation was derived directly from an underlying microscopic model irrespective of environmental memory time and coupling strength. Several physically interesting models have been solved using the non-Markovian QSD approach ranging from one qubit [19], two-qubit [20], N-qubit systems [21], to multi-level systems [22, 23]. Moreover, recent research has extended the theory of QSD for a bosonic bath to an open system coupled to a fermionic bath [24, 25]. Furthermore, as a computing tool in real applications, the QSD approach also showed its potential value in many other interesting problems including a theory of precise measurement [27], quantum control dynamics [28], and quantum biology [29].

The non-Markovian QSD equation has been widely applied to discrete-variable systems including qubits, qutrits, spins etc. However, it is also important to consider the non-Markovian QSD approach for multiple continuous-variable (CV) systems since CV systems are known to have many unique features leading to advantages in its application in quantum information processing (QIP) such as quantum teleportation, quantum error-correction codes, dense coding, and quantum computing [30]. Furthermore, CV systems are an integral part of quantum dissipative systems, non-equilibrium statistical physics and theory of quantum decoherence [3].

The purpose of this paper is to derive the non-Markovian QSD equation and the corresponding master equation for a coupled N-cavity model. This is a very good example of using the QSD technique to solve a multiple CV quantum open system. It should be noted that the coupled N-cavity model itself is an important model in QIP due to its relevance to the quantum communication network schemes [31]. As simple cases of the coupled N-cavity model, single-cavity and coupled two-cavity have been widely studied and the exact master equations are derived by using the path integral approach [7, 9, 10]. We extend the two-cavity case into a multi-cavity case and show that the QSD technique is a very useful tool in dealing with the N-cavity model. Then, based on the non-Markoivan QSD equations and master equations, we also study several interesting phenomena concerning quantum coherence transfer assisted by the environment-memory. Specifically, in the two-cavity example, we show that the memory-effect of highly non-Markovian environment can be used to transfer the so-called Schrödinger cat-like state between the two cavities. This phenomenon exists only in the strongly non-Markovian environment, so it may be called the memory-assisted state transfer. Furthermore, in the case of the three coupled cavities model, we study the cat-like state transfer and entanglement transfer under different boundary conditions, and show that the dynamics of the system is strongly dependent on the boundary conditions.

This paper is organized as follows: In Sec. II, we derive the exact QSD equation and the corresponding master equation for coupled N-cavity model. The finite temper-

---

*Email:xzhao1@stevens.edu
†Email:Ting.Yu@stevens.edu

ature case is subsequently discussed in Sec. III. Then, in Sec. IV, we study a special example, the coupled two-cavity case. We show that the cat-like state can be transferred from cavity 1 to cavity 2 without any direct couplings. Furthermore, we show that the cat-like state transfer can only be observed in a highly non-Markovian environment. Next, in Sec. V, the coupled three-cavity example is considered. We show that the quantum dynamics (cat-like state transfer or entanglement transfer) will be significantly modified by the boundary conditions. Finally, we conclude in Sec. VI.

II. COUPLED N-CAVITY MODEL: ZERO-TEMPERATURE BATH

We consider a coupled cavity array interacting with a common bosonic bath. The total Hamiltonian of the system plus environment can be written as,

$$H_{\rm tot} = H_{\rm s} + H_{\rm b} + H_{\rm int}, \tag{1}$$

where the Hamiltonian of the system is described by

$$H_{\rm s} = \sum_{i=1}^{N} \omega_i a_i^\dagger a_i + \sum_{i=1}^{N} \lambda_i (a_i^\dagger a_{i+1} + a_i a_{i+1}^\dagger), \tag{2}$$

where ω_i $(i = 1, 2, ...N)$ are the frequencies for N cavities, and λ_i are the coupling constants between ith cavity and $(i+1)th$ cavity. We consider two types of boundary conditions for this model. The first one is the periodical boundary condition (PBC), which means that N cavities form a closed loop (i.e., $a_{N+1} = a_1$). The other one is the open boundary condition (OBC). Namely, the cavity chain has two open ends. We assume that all the cavities are coupled to a common bath, which is composed of a set of harmonic oscillators b_j, $b_j^\dagger$.

$$H_{\rm b} = \sum_j \nu_j b_j^\dagger b_j. \tag{3}$$

The interaction Hamiltonian between the system and the environment is

$$H_{\rm int} = \sum_{i,j} (k_{ij} b_j^\dagger a_i + k_{ij}^* b_j a_i^\dagger). \tag{4}$$

For simplicity, we assume that each $k_{ij} = g_j l_i$ is the product of two coupling constants g_j and l_i, then

$$H_{\rm int} = \sum_j (g_j b_j^\dagger L + g_j^* b_j L^\dagger), \tag{5}$$

where the Lindblad operator $L = \sum_{i=1}^{N} l_i a_i$. In the following derivation, we set all the coefficients $l_i = 1$ for simplicity.

The formal Diósi-Gisin-Strunz equation can be written as [18],

$$\frac{\partial}{\partial t}\psi_t = -iH_{\rm s}\psi_t + L z_t^* \psi_t - L^\dagger \int_0^t ds \alpha(t,s) \frac{\delta \psi_t}{\delta z_s^*}, \tag{6}$$

where $\alpha(t,s) = \sum_j |g_j|^2 e^{-i\nu_j (t-s)}$ is the bath correlation function at zero temperature $(T = 0)$, and $z_t^* = -i\sum_j g_j z_j^* e^{i\nu_j t}$ is a complex Gaussian process satisfying $M[z_t] = M[z_t z_s] = 0$, $M[z_t^* z_s] = \alpha(t,s)$. Here $M[\cdot]$ denotes the statistical average over the classical noise z_t^*. The reduced density operator of the system may be recovered by the statistical average from the quantum trajectories $\psi_t(z^*)$: $\rho_t = M[|\psi_t(z^*)\rangle\langle\psi_t(z^*)|] = \int \frac{dz^2}{\pi} e^{-|z|^2} |\psi_t(z^*)\rangle\langle\psi_t(z^*)|$.

In order to establish a time-local non-Markovian QSD equation, we have to replace the functional derivative $\frac{\delta\psi_t}{\delta z_s^*}$ by a time-local operator $O(t,s,z^*)$,

$$\frac{\delta \psi_t(z^*)}{\delta z_s^*} = O(t,s,z^*)\psi_t(z^*). \tag{7}$$

We will use the shorthand notation: $O = O(t,s,z^*)$ if no confusion arises. According to the consistency condition $\frac{\delta}{\delta z_s^*}\frac{\partial}{\partial t}\psi_t(z^*) = \frac{\partial}{\partial t}\frac{\delta}{\delta z_s^*}\psi_t(z^*)$, the operator O can be shown to satisfy the following dynamic equation,

$$\frac{\partial}{\partial t} O = [-iH_{\rm s} + L z_t^* - L^\dagger \bar{O}, O] - L^\dagger \frac{\delta}{\delta z_s^*}\bar{O}, \tag{8}$$

where $\bar{O}$ is defined as $\bar{O}(t,s,z^*) = \int_0^t ds \alpha(t,s) O(t,s,z^*) ds$. The initial condition is given by

$$O(t,t,z^*) = L = \sum_{i=1}^{N} a_i. \tag{9}$$

From Eq. (8), the exact O operator of this model can be derived as,

$$O(t,s) = \sum_{i=1}^{N} f_i(t,s) a_i. \tag{10}$$

Note that the O operator for this coupled cavity array does not contain the noise z_t^*. Substituting Eq. (10) into Eq. (8), we can show that the coefficient functions $f_i(t,s)$ satisfy the following equations,

$$\begin{aligned} \frac{\partial}{\partial t} f_i(t,s) &= i\omega_i f_i(t,s) + i[\lambda_i f_{i+1}(t,s) + \lambda_{i-1} f_{i-1}(t,s)] \\ &\quad + \sum_{j=1}^{N} f_j(t,s) F_i(t), \end{aligned} \tag{11}$$

where $F_i(t) = \int_0^t ds \alpha(t,s) f_i(t,s)$ $(i = 1, 2, .., N)$, with the initial conditions

$$f_i(t = s, s) = 1. \tag{12}$$

With the exact O operator given by Eq. (10), we can derive the exact master equation for the system from the exact QSD equation [19],

$$\begin{aligned} \frac{d}{dt}\rho &= -i[H_{\rm s}, \rho] + [L, \rho \bar{O}^\dagger(t)] + [\bar{O}(t)\rho, L^\dagger] \\ &= -i[H_{\rm s}, \rho] + \{ \sum_{i,j=1}^{N} F_j^*(t)[a_i, \rho a_j^\dagger] + \text{h.c.} \}. \end{aligned} \tag{13}$$

This exact master equation is derived in a very simple manner due to the fact that O operator is noise-independent. However, when the initial state of the bath is a finite-temperature thermal state, the operator O will not be noise free, and the derivation of the exact master equation becomes more involved as to be shown in the next section.

III. COUPLED N-CAVITY MODEL: FINITE-TEMPERATURE BATH

In the case of the finite-temperature bath, we can introduce a fictitious bath and transform the finite temperature case into an effective zero temperature model [18, 32]. The non-Markovian QSD equation for the finite-temperature case is given by

$$\frac{\partial}{\partial t}\psi_t = (-iH_{\rm s} + Lz_t^* + L^\dagger w_t^* - L^\dagger\bar{O}_1 - L\bar{O}_2)\psi_t, \quad (14)$$

where $H_{\rm s}$ is given in Eq. (2), and $z_t^* = -i\sum_i f_i z_i^* e^{i\nu_i t}$, $w_t^* = -i\sum_i h_i^* w_i^* e^{-i\nu_i t}$ are two statistically independent Gaussian noises with $f_i = \sqrt{\bar{n}_i+1}g_i$ and $h_i = \sqrt{\bar{n}_i}g_i$ as the effective coupling constants. Here $\bar{n}_i = \frac{1}{\exp(\hbar\nu_i/k_BT)-1}$. The two Gaussian noises satisfy the following relations,

$$M[z_t] = M[z_t z_s] = 0, \quad M[z_t^* z_s] = \alpha_1(t,s), \quad (15)$$

$$M[w_t] = M[w_t w_s] = 0, \quad M[w_t^* w_s] = \alpha_2(t,s). \quad (16)$$

where $\alpha_1(t,s) = \sum_i |f_i|^2 e^{-i\nu_i(t-s)}$ and $\alpha_2(t,s) = \sum_i |h_i|^2 e^{i\nu_i(t-s)}$ are correlation functions for the two effective baths. For the finite-temperature case, the two O operators are defined as

$$O_1(t,s,z^*,w^*)\psi_t = \frac{\delta\psi_t(z^*,w^*)}{\delta z_s^*}, \quad (17)$$

$$O_2(t,s,z^*,w^*)\psi_t = \frac{\delta\psi_t(z^*,w^*)}{\delta w_s^*}, \quad (18)$$

satisfying the following equations

$$\frac{\partial}{\partial t}O_1 = [-iH_{\rm s} + Lz_t^* + L^\dagger w_t^* - L^\dagger\bar{O}_1 - L\bar{O}_2, O_1] - L^\dagger\frac{\delta}{\delta z_s^*}\bar{O}_1 - L\frac{\delta}{\delta z_s^*}\bar{O}_2, \quad (19)$$

$$\frac{\partial}{\partial t}O_2 = [-iH_{\rm s} + Lz_t^* + L^\dagger w_t^* - L^\dagger\bar{O}_1 - L\bar{O}_2, O_2] - L^\dagger\frac{\delta}{\delta w_s^*}\bar{O}_1 - L\frac{\delta}{\delta w_s^*}\bar{O}_2, \quad (20)$$

where $\bar{O}_i = \int_0^t \alpha_i(t,s)O_i(t,s,z^*,w^*)ds$ $(i=1,2)$, According to Eq. (19-20), the exact O operators for the N-cavity model can be determined as

$$O_1(t,s,w^*) = \sum_{i=1}^N p_i(t,s)a_i + \int_0^t q(t,s,s')w_{s'}^* ds', \quad (21)$$

$$O_2(t,s,z^*) = \sum_{i=1}^N x_i(t,s)a_i^\dagger + \int_0^t y(t,s,s')z_{s'}^* ds', \quad (22)$$

while the coefficients satisfying the following equations

$$\begin{aligned}\frac{\partial}{\partial t}p_i(t,s) &= i\omega_i p_i(t,s) + i[\lambda_i p_{i+1}(t,s) + \lambda_{i-1}p_{i-1}(t,s)] \\ &+ \sum_{j=1}^N p_j(t,s)P_i(t) + \sum_{j=1}^N p_j(t,s)X_j(t) - Y(t,s),\end{aligned} \quad (23)$$

$$\begin{aligned}\frac{\partial}{\partial t}x_i(t,s) &= -i\omega_i x_i(t,s) - i[\lambda_i x_{i+1}(t,s) + \lambda_{i-1}x_{i-1}(t,s)] \\ &- \sum_{j=1}^N x_j(t,s)P_j(t) - \sum_{j=1}^N x_j(t,s)X_i(t) - Q(t,s),\end{aligned} \quad (24)$$

$$\frac{\partial}{\partial t}q(t,s,s') = \sum_{j=1}^N p_j(t,s)Q(t,s'), \quad (25)$$

$$\frac{\partial}{\partial t}y(t,s,s') = -\sum_{j=1}^N x_j(t,s)Y(t,s'), \quad (26)$$

where

$$P_i(t) = \int_0^t \alpha_1(t,s)p_i(t,s)ds, \quad (27)$$

$$Q(t,s') = \int_0^t \alpha_1(t,s)q(t,s,s')ds, \quad (28)$$

$$X_i(t) = \int_0^t \alpha_2(t,s)x_i(t,s)ds, \quad (29)$$

$$Y(t,s') = \int_0^t \alpha_2(t,s)y(t,s,s')ds, \quad (30)$$

with the initial conditions

$$p_i(t,t) = 1, \quad (31)$$

$$x_i(t,t) = 1, \quad (32)$$

$$q(t,t,s') = 0, \quad (33)$$

$$y(t,t,s') = 0, \quad (34)$$

$$q(t,s,t) = -\sum_{j=1}^N p_j(t,s), \quad (35)$$

$$y(t,s,t) = \sum_{j=1}^N x_j(t,s). \quad (36)$$

Since the O operators in Eq. (21-22) are noise-dependent, the exact master equation cannot be obtained directly from Eq. (13). We have to derive the new master equation in the finite temperature case. Using the so-called

Heisenberg approach [26], the exact master equation is derived as

$$\begin{aligned}\frac{\partial}{\partial t}\rho &= -i[H_{\rm s},\rho] \\ &\quad + \{C_1^*(t)[\sum_i a_i, \sum_j \rho a_j^\dagger] + C_2^*(t)[\sum_i a_i, \sum_j a_j^\dagger \rho] \\ &\quad + C_3^*(t)[\sum_i a_i^\dagger, \sum_j \rho a_j] + C_4^*(t)[\sum_i a_i^\dagger, \sum_j a_j \rho] \\ &\quad + \text{h.c.}\}, \qquad (37)\end{aligned}$$

where the coefficients $C_i(t)$ $(i = 1, 2, 3, 4)$ are given by

$$\begin{aligned} C_i(t) &= \int_0^t \alpha_1(t,s) c_i(t,s) ds \quad (i = 1, 2), \\ C_j(t) &= \int_0^t \alpha_2(t,s) c_j(t,s) ds \quad (j = 3, 4), \qquad (38)\end{aligned}$$

and the details of the derivation will be shown in Appendix A.

This is the general form of the master equation for N-cavity model in the finite temperature case. This model serves as a good example showing that the QSD approach is a useful tool in deriving the non-Markovian master equation in both zero temperature and finite temperature case. In the following two sections, we will discuss several examples with $N = 2$ and $N = 3$.

IV. TWO COUPLED CAVITIES

In this and the next sections, we will show several interesting numerical results based on the simple examples $N = 2$ and $N = 3$. The non-Markovian QSD equation for the model consisting of one cavity was discussed in [19], here we start from the case of two coupled cavities. For the case with $N = 2$, the system Hamiltonian and the Lindblad operator are given by,

$$H_{\rm s} = \omega_1 a_1^\dagger a_1 + \omega_2 a_2^\dagger a_2 + \lambda(a_1^\dagger a_2 + a_1 a_2^\dagger), \qquad (39)$$

$$L = a_1 + a_2. \qquad (40)$$

According to the general QSD equation for N-cavity model in Sec. II, the non-Markovian QSD equation for the coupled two-cavity model is

$$\frac{\partial}{\partial t}\psi_t = (-iH_{\rm s} + L z_t^* - L^\dagger \bar{O})\psi_t, \qquad (41)$$

where $\bar{O}(t) = F_1(t)a_1 + F_2(t)a_2$, the coefficients are determined by Eq. (11). From Eq. (13), the corresponding master equation for the coupled two-cavity model is

$$\begin{aligned}\frac{\partial}{\partial t}\rho &= -i\omega_1(a_1^\dagger a_1 \rho - \rho a_1^\dagger a_1) - i\omega_2(a_2^\dagger a_2 \rho - \rho a_2^\dagger a_2) \\ &\quad - i\lambda(a_1^\dagger a_2 \rho - \rho a_1^\dagger a_2 + a_2^\dagger a_1 \rho - \rho a_2^\dagger a_1) \\ &\quad + \{F_1(t)(a_1 \rho a_1^\dagger - a_1^\dagger a_1 \rho + a_1 \rho a_2^\dagger - a_2^\dagger a_1 \rho) \\ &\quad + F_2(t)(a_2 \rho a_1^\dagger - a_1^\dagger a_2 \rho + a_2 \rho a_2^\dagger - a_2^\dagger a_2 \rho) + \text{h.c.}\}. \qquad (42)\end{aligned}$$



FIG. 1: (Color online) Wigner functions for the two cavities are plotted at $\omega t = 0$, $\omega t = 11$, $\omega t = 22$, and $\omega t = 33$. The initial state is chosen as the cat-state $1/\sqrt{Z}(|1\rangle + |-1\rangle) \otimes |0\rangle$. The parameters are $\omega_1 = \omega_2 = \omega = 1$, $\lambda = 0$, $\gamma = 0.1$.

Note that our derivations of the QSD and master equation are independent on the choice of the environment correlation functions. For simplicity, all the numerical simulations using the QSD equation are based on the Ornstein-Ulenbeck noise, which is defined by $\alpha(t,s) = \frac{\gamma}{2} e^{-\gamma|t-s|}$. The Ornstein-Ulenbeck process recovers the Markov limit when $\gamma \to \infty$, $\alpha(t,s) \to \delta(t,s)$. This property will allow us to observe the crossover properties from non-Markovian to Markov regimes. A small γ typically represents a non-Markovian noise with the finite memory time.

For two coupled cavities without interacting with an environment, one can easily see that the phenomenon of cat-like state transfer due to the mutual coupling between the two cavities. In fact, if we choose the initial state of the first cavity as the Schrödinger's cat-like state $|\psi_{\rm cat}\rangle = 1/\sqrt{Z}(|\alpha\rangle + |-\alpha\rangle)$ (Z is a normalization factor), while the second cavity is in the vacuum state, i.e., $|\psi(0)\rangle = 1/\sqrt{Z}(|\alpha\rangle + |-\alpha\rangle)_1 \otimes |0\rangle_2$, the cat-like state will be transferred to the second cavity from the first cavity after time t_c: $|\psi(t = t_c)\rangle = |0\rangle_1 \otimes 1/\sqrt{Z}(|\alpha\rangle + |-\alpha\rangle)_2$. This result can be simply proved by solving the corresponding Fokker-Planck equation [33].

Using the exact QSD equation and master equation, we study several interesting properties of non-Markovian dynamics of the multiple cavity model. We find that the non-Markovian common bath can induce a mutual correlation between two cavities indirectly, and make the cat-like state hop from one cavity to another even the two cavities are not interacting to each other. Notably, this phenomenon of the memory-assisted transfer can be only observed in a strongly non-Markovian environment. It is easy to show that the Markov environment will quickly suppress the possibility of the state transfer due to the irreversible energy dissipation to the environment. In Fig. 1, we plot the Wigner functions for the two cavities at different time points. The regions with negative values reflect the quantum interference due to the coherent superposition of two wave packets. In a strongly non-Markovian case ($\gamma = 0.1$), the cat-like state can be transferred to the second cavity from the first cavity without any direct coupling between two cavities (i.e., $\lambda = 0$). As a comparison, it is noted that when the environment is in a Markov regime (γ is large), the high fidelity transfer cannot happen (the Markov case is not plotted in the figure). Since the dissipative process is dominant in the Markov limit, the decoherence process suppresses the transfer process swiftly. Therefore, the cat-like state may be destroyed by dissipation long before it can hop to the second cavity. The numerical results show how a non-Markovian environment affects the quantum state evolution.

V. THREE COUPLED CAVITIES

A more complicated example is the case of three coupled cavities. For $N = 3$, the system Hamiltonian can be written as

$$\begin{aligned} H_s = {} & \omega_1 a_1^\dagger a_1 + \omega_2 a_2^\dagger a_2 + \omega_3 a_3^\dagger a_3 + \lambda_1 (a_1^\dagger a_2 + a_1 a_2^\dagger) \\ & + \lambda_2 (a_2^\dagger a_3 + a_2 a_3^\dagger) + \lambda_3 (a_1^\dagger a_3 + a_1 a_3^\dagger), \end{aligned} \tag{43}$$

and the Lindblad operator is given by,

$$L = a_1 + a_2 + a_3. \tag{44}$$

This model allows us to consider two types of boundary conditions. In the case of the open boundary condition (OBC), we have $\lambda_1 = \lambda_2 = \lambda$, $\lambda_3 = 0$. In contrast, in the case of the periodic boundary condition (PBC), we have following constraints: $\lambda_1 = \lambda_2 = \lambda_3 = \lambda$. According to the QSD equation for the general N-cavity model discussed in Sec. II, the exact QSD equation is given by

$$\frac{\partial}{\partial t}\psi_t = (-iH_s + L z_t^* - L^\dagger \bar{O})\psi_t, \tag{45}$$

where $\bar{O} = F_1(t)a_1 + F_2(t)a_2 + F_3(t)a_3$, and the coefficients are determined by Eq. (11). Consequently, the corresponding exact master equation is given by,

$$\begin{aligned} \frac{\partial}{\partial t}\rho = {} & -i\omega_1(a_1^\dagger a_1 \rho - \rho a_1^\dagger a_1) - i\omega_2(a_2^\dagger a_2 \rho - \rho a_2^\dagger a_2) \\ & - i\omega_3(a_3^\dagger a_3 \rho - \rho a_3^\dagger a_3) \\ & - i\lambda_1(a_1^\dagger a_2 \rho - \rho a_1^\dagger a_2 + a_2^\dagger a_1 \rho - \rho a_2^\dagger a_1) \\ & - i\lambda_2(a_2^\dagger a_3 \rho - \rho a_2^\dagger a_3 + a_3^\dagger a_2 \rho - \rho a_3^\dagger a_2) \\ & - i\lambda_3(a_1^\dagger a_3 \rho - \rho a_1^\dagger a_3 + a_3^\dagger a_1 \rho - \rho a_3^\dagger a_1) \\ & + \{F_1^*(t)(a_1 \rho a_1^\dagger - \rho a_1^\dagger a_1 + a_2 \rho a_1^\dagger \\ & - \rho a_1^\dagger a_2 + a_3 \rho a_1^\dagger - \rho a_1^\dagger a_3) \\ & + F_2^*(t)(a_2 \rho a_2^\dagger - \rho a_2^\dagger a_2 + a_1 \rho a_2^\dagger \\ & - \rho a_2^\dagger a_1 + a_3 \rho a_2^\dagger - \rho a_2^\dagger a_3) \\ & + F_3^*(t)(a_3 \rho a_3^\dagger - \rho a_3^\dagger a_3 + a_1 \rho a_3^\dagger \\ & - \rho a_3^\dagger a_1 + a_2 \rho a_3^\dagger - \rho a_3^\dagger a_2) + \text{h.c.}\}. \end{aligned} \tag{46}$$



FIG. 2: (Color online) Time evolution of fidelities. (a) is for the case of PBC, and (b) is for the case of OBC. The red (solid), green (dashed), blue (dash-dotted) lines are the fidelities $\mathcal{F}^{(1)}$, $\mathcal{F}^{(2)}$, and $\mathcal{F}^{(3)}$ for cavities 1, 2, 3, respectively. The parameters are $\omega_1 = \omega_2 = \omega_3 = \omega = 1$, $\lambda = 1$, $\gamma = 0.2$.



FIG. 3: (Color online) Wigner functions for three cavities in the case of OBC at different time points. The initial state is chosen as $1/\sqrt{Z}(|1\rangle + |-1\rangle) \otimes |0\rangle \otimes |0\rangle$. The parameters are $\omega_1 = \omega_2 = \omega_3 = \omega = 1$, $\lambda_1 = \lambda_2 = 1$, $\lambda_3 = 0$, $\gamma = 0.2$.

The above exact QSD equation and master equation will serve as a powerful tool in our study on the properties of the non-Markovian dynamics under two different boundary conditions.



FIG. 4: (Color online) Wigner functions for three cavities in the case of PBC at different time points. The initial state is chosen as $1/\sqrt{Z}(|1\rangle + |{-1}\rangle) \otimes |0\rangle \otimes |0\rangle$. The parameters are $\omega_1 = \omega_2 = \omega_3 = \omega = 1$, $\lambda_1 = \lambda_2 = \lambda_3 = 1$, $\gamma = 0.2$.



FIG. 5: (Color online) Time evolution of the negativity for all the three modes. The red (solid), green (dashed) and blue (short-dashed) lines are entanglement (negativity) between cavities 1-2, 1-3 and 2-3, respectively. The initial state is chosen as $|\psi(0)\rangle = \frac{1}{\sqrt{2}}(|100\rangle + |010\rangle)$. The parameters are $\omega_1 = \omega_2 = \omega_3 = \omega = 1$, $\lambda = 1$, $\gamma = 0.2$.

A. Cat-like state transfer under different boundary conditions

We have already showed that the cat-like state $1/\sqrt{Z}(|\alpha\rangle + |-\alpha\rangle)$ can be transferred from one cavity to another both analytically and numerically in the case of $N = 2$. However, the cat-like state transfer can also be realized in the $N = 3$ case. Moreover, the model of three coupled cavities allows us to consider the two different boundary conditions that will profoundly affect the effect of state transfer. In Fig. 2, fidelities are plotted for three cavities under two different boundary conditions. The fidelities are calculated as $\mathcal{F}^{(i)} = \max_{\{\theta\}}[\langle\psi_{\text{cat}}^{(R)}(\theta)|\rho^{(i)}(t)|\psi_{\text{cat}}^{(R)}(\theta)\rangle]$, where $|\psi_{\text{cat}}^{(R)}(\theta)\rangle = 1/\sqrt{Z}(|\alpha e^{-i\theta}\rangle + |-\alpha e^{-i\theta}\rangle)$ is the rotated cat-like state, $\rho^{(i)}(t)$ is the reduced density matrix for the cavity i, and $\mathcal{F}^{(i)}$ is the fidelity for the cavity i. Since the cat state may rotate in time evolution no matter with or without an environment, so all the rotated cat-like state $|\psi_{\text{cat}}^{(R)}(\theta)\rangle$ for any given θ can be considered as a transfered cat-like state. Therefore, we calculate the maximum fidelity between $\rho^{(i)}(t)$ and $|\psi_{cat}^{(R)}(\theta)\rangle$ for all possible θ. From the curves of fidelities in Fig. 2, it is shown that in the case of PBC, the fidelity will increase and reach to 1 only in the first cavity. In the case of OBC, the fidelity of the third cavity can be close to 1 at $\omega t \approx 2.2$, which means that the cat-like state is transferred into the third cavity. It is worthwhile to point out that, in the case of OBC, the cat-like state will only be transferred from cavity 1 to cavity 3. It means that the state of the cavity 2 will never be a cat state. Intuitively, in the case of OBC, two cavities at the end points of a chain are symmetric while the middle cavity is in a unique position. This is similar to the classical model of a mechanical oscillator in a string, the superposition of the oscillation forms a standing wave due to the boundary conditions, so that the middle point on the string is just a “node”, whose amplitude is always zero. In contrast, for the case of PBC, the cat-like state never transfer into other cavities since the other two cavities are in the symmetric positions. The state has an equal probability to be one of the two cavities, so the states of two cavity cannot be the cat state at the same time. Therefore, one will expect that the cat state will experience a decay-revival process in the initial cavity. To examine this more explicitly, in Fig. 3 and Fig. 4, we plot the Wigner functions for each cavity in the three-cavity model with OBC and PBC. The time evolution of the Wigner functions shows that the cat-like state is able to transfer into cavity 3 at $\omega t = 2.2$ in OBC case. However, in the case of PBC,one only observe the decay and revival of the cat-like state in the first cavity (initial cavity) at $\omega t = 2.04$. Of course, the cat state will disappear completely in a long-time limit due to the dissipation effect.

B. Entanglement transfer under different boundary conditions

As another interesting phenomenon, let us consider the entanglement transfer in the cavity array. We choose the initial state of the three-cavity chain as $|\psi(0)\rangle =$

$\frac{1}{\sqrt{2}}(|100\rangle + |010\rangle)$ (in Fock basis), where cavity 1 and 2 are entangled, but the combined cavity 1 and 2 are separable from cavity 3. In the OBC case, the initial entanglement between cavity 1 and 2 will transfer to the combined system of cavity 2 and 3. Specifically, in Fig. 5 (OBC case), the entanglement between cavity 2 and 3 will reach the maximum value at $\omega t \approx 2.2$. At the same time point, the red line and green line in Fig. 5 show that cavity 1 become separable from both cavity 2 and cavity 3. As a comparison, we see that, in the PBC case, the entanglement between cavity 2 and 3 never reach 1. However, PBC case is shown to be more helpful to the entanglement preservation. For example, the loss of entanglement degree in cavity 1 and 2 at $\omega t \approx 4$ is less than that in the case of OBC.

VI. CONCLUSION

In this paper, we have derived the time-local non-Markovian QSD for the coupled N-cavity model embedded in a common bath without any approximation. Using the non-Markovian quantum trajectory approach, we have successfully derived the exact master equation for the N coupled cavity model in both the zero and finite-temperature bath cases. Based on these exact evolution equations, we studied several interesting phenomena in the dynamic evolution of the two-cavity and three-cavity cases. We studied the cat-like state transfer in the coupled cavity array, and showed that a highly non-Markovian noise can enhance the cat-like state transfer (called the memory-assisted cat state transfer). In the three-cavity example, we also studied the effect of boundary conditions on the cat-like state transfer and the entanglement transfer. These examples are used to show that the non-Markovian QSD approach is a highly valuable tool in characterizing the dynamics of a CV system. It is important to note that the approach can be directly extended to a generic multipartite CV system with an arbitrary number of cavities.

Acknowledgments

We acknowledge grant support from the NSF PHY-0925174, DOD/AF/AFOSR No. FA9550-12-1-0001, The NBRPC No. 2009CB929300, the NSFC Nos. 91121015 and the MOE No. B06011.

Appendix A: Deriving the non-Markovian master equation in the finite temperature case

In this section, we will provide details of deriving the finite temperature master equations from the exact QSD equations. We only consider the PBC case, and notice that the OBC case can be treated in a similar way. For simplicity, we assume that all the cavities are identical (i.e., $\omega_i = \omega$, $\lambda_i = \lambda$). Following the Heisenberg approach [26, 32], one can deal with the derivation of the exact master equation in the general case.

In the finite temperature case [18, 32], the total Hamiltonian in the interaction picture is given by

$$H_{\mathrm{tot}}^{(I)}(t) = H_{\mathrm{s}} + \sum_i (f_i^* e^{-i\nu_i t} L^\dagger d_i + f_i e^{i\nu_i t} L d_i^\dagger) + \sum_i h_i^* e^{-i\nu_i t} L^\dagger e_i^\dagger + h_i e^{i\nu_i t} L e_i), \tag{A1}$$

where the Hamiltonian of the system (coupled N-cavity) can be written as

$$H_{\mathrm{s}} = \omega \sum_{i=1}^{N} a_i^\dagger a_i + \lambda \sum_{i=1}^{N} (a_i^\dagger a_{i+1} + a_{i+1}^\dagger a_i). \tag{A2}$$

In the Heisenberg picture, the evolution for any operator is given by, for example,

$$\frac{\partial}{\partial t} d_i(t) = -iU_t^{-1}[d_i, H_{\mathrm{tot}}^{(I)}(t)]U_t = -i f_i e^{i\nu_i t} L\,(t)\,. \tag{A3}$$

Integrating both sides of the equation and noticing the initial condition $d_i(0) = d_i$, we obtain

$$d_i(t) = d_i - i f_i \int_0^t L(s) e^{i\nu_i s} ds, \tag{A4}$$

similarly, we also have

$$e_i(t) = d_i - i h_i^* \int_0^t L^\dagger(s) e^{-i\nu_i s} ds, \tag{A5}$$

$$d_i^\dagger(s) = d_i^\dagger(t) - i f_i^* \int_s^t L^\dagger(s') e^{-i\nu_i s'} ds', \tag{A6}$$

$$e_i^\dagger(s) = e_i^\dagger(t) - i h_i \int_s^t L(s') e^{i\nu_i s'} ds', \tag{A7}$$

For a single trajectory, the time evolution of the state of the system is governed by the stochastic propagator of the system $G(t, z^*, w^*) = \langle z|\langle w|U_t|0\rangle$ as

$$|\psi\,(t, z^*, w^*)\rangle = G(t, z^*, w^*)|\psi_{\mathrm{s}}\,(0)\rangle. \tag{A8}$$

Applying Schrödinger equation to the whole system and noticing Eq. (A4-A7), we will find $G(t, z^*, w^*)$ satisfies an equation similar to the QSD equation (14),

$$\begin{aligned}\frac{\partial}{\partial t} & G(t, z^*, w^*) \\ &= -iH_s G + L z_t^* G - L^\dagger \int_0^t ds \alpha_1(t,s) \langle z|\langle w|U_t L(s)|0\rangle \\ &+ L^\dagger w_t^* G - L \int_0^t ds \alpha_2(t,s) \langle z|\langle w|U_t L^\dagger(s)|0\rangle,\end{aligned} \tag{A9}$$

where $\alpha_1(t,s) = \sum_i |f_i|^2 e^{-i\nu_i(t-s)}$, $\alpha_2(t,s) = \sum_i |h_i|^2 e^{i\nu_i(t-s)}$. From this dynamic equation, we can also define the O operators from the stochastic propagator

$$\langle z|\langle w|U_t L(s)|0\rangle = O_1(t,s,z^*,w^*)G(t,z^*,w^*), \tag{A10}$$

$$\langle z|\langle w|U_t L^\dagger(s)|0\rangle = O_2(t,s,z^*,w^*)G(t,z^*,w^*). \tag{A11}$$

Instead of deriving the non-linear differential equations of O operator with respect to t such as Eq. (23-26), we will derive another set of differential equations with respect to the initial value s, and it will be showed later that these equations are linear. Taking the derivative with respect to s of Eq. (A10), one gets,

$$\begin{aligned}
&\frac{\partial}{\partial s}\langle z|\langle w|U_t L(s)|0\rangle \\
&= \sum_i \langle z|\langle w|U_t(-iU_s^{-1}[a_i, H_{\rm tot}^{(I)}(s)]U_s)|0\rangle \\
&= -i\sum_i \omega\langle z|\langle w|U_t a_i(s)|0\rangle \\
&- i\sum_i \lambda\langle z|\langle w|U_t[a_{i+1}(s)+a_{i-1}(s)]|0\rangle \\
&- iN\langle z|\langle w|U_t \sum_i f_i^* e^{-i\nu_i s} d_i(s)|0\rangle \\
&- iN\langle z|\langle w|U_t \sum_i h_i^* e^{-i\nu_i s} e_i^\dagger(s)|0\rangle,
\end{aligned} \tag{A12}$$

and noticing Eq. (A4-A7),

$$\begin{aligned}
&\frac{\partial}{\partial s}\langle z|\langle w|U_t L(s)|0\rangle \\
&= -i\omega\langle z|\langle w|U_t L(s)|0\rangle - 2i\lambda\langle z|\langle w|U_t L(s)|0\rangle \\
&- N\int_0^s ds'\alpha_1(s,s')\langle z|\langle w|U_t L(s^{'})|0\rangle + Nw_s^*\langle z|\langle w|U_t|0\rangle \\
&- N\int_s^t ds'\alpha_2(s',s)\langle z|\langle w|U_t L(s')|0\rangle.
\end{aligned} \tag{A13}$$

So,

$$\begin{aligned}
\frac{\partial}{\partial s}O_1 &= -i\omega O_1 - 2i\lambda O_1 \\
&\quad - N\int_0^s ds'\alpha_1(s,s')O_1(t,s',z^*,w^*) \\
&\quad + Nw_s^* - N\int_s^t ds'\alpha_2(s',s)O_1(t,s',z^*,w^*).
\end{aligned} \tag{A14}$$

Similarly, we can obtain

$$\begin{aligned}
\frac{\partial}{\partial s}O_2 &= i\omega O_2 + 2i\lambda O_2 \\
&\quad + N\int_0^s ds'\alpha_2(s,s')O_2(t,s',z^*,w^*) \\
&\quad - Nz_s^* + N\int_s^t ds'\alpha_1(s',s)O_2(t,s',z^*,w^*).
\end{aligned} \tag{A15}$$

Thus, we have obtained the differential equations for O_1 and O_2 with respect to s.

Recall the master equation derived in Ref. [32],

$$\begin{aligned}
\frac{\partial}{\partial t}\rho = -i[H_s,\rho] &+ [L, M\{P_t\bar{O}_1^\dagger\}] - [L^\dagger, M\{\bar{O}_1 P_t\}] \\
&+ [L^\dagger, M\{P_t\bar{O}_2^\dagger\}] - [L, M\{\bar{O}_2 P_t\}].
\end{aligned} \tag{A16}$$

Define

$$R_1(t,s) = M\{O_1 P_t\},\ R_1^\dagger(t,s) = M\{P_t O_1^\dagger\}, \tag{A17}$$

$$R_2(t,s) = M\{O_2 P_t\},\ R_2^\dagger(t,s) = M\{P_t O_2^\dagger\}. \tag{A18}$$

then, from Eq. (A14-A15), we will obtain,

$$\begin{aligned}
\frac{\partial}{\partial s}R_1 = -i\omega R_1 &- 2i\lambda R_1 - N\int_0^s ds'\alpha_1(s,s')R_1(t,s') \\
&+ N\int_0^t ds'\alpha_2^*(s,s')R_2^\dagger(t,s') \\
&- N\int_s^t ds'\alpha_2(s',s)R_1(t,s'),
\end{aligned} \tag{A19}$$

with the initial condition

$$R_1(t,t) = L\rho. \tag{A20}$$

In the derivation above, the Novikov theorem

$$\begin{aligned}
M\{w_s^* P_t\} &= \int_0^t ds'\alpha_2^*(s,s')M\{P_t O_2^\dagger(t,s',z^*,w^*)\} \\
&= \int_0^t ds'\alpha_2^*(s,s')R_2^\dagger(t,s'),
\end{aligned} \tag{A21}$$

is used.

Similarly, equation for R_2 can be derived as

$$\begin{aligned}
\frac{\partial}{\partial s}R_2 = i\omega R_2 &+ 2i\lambda R_2 + N\int_0^s ds'\alpha_2(s,s')R_2(t,s') \\
&- N\int_0^t ds'\alpha_1^*(s,s')R_1^\dagger(t,s') \\
&+ N\int_s^t ds'\alpha_1(s',s)R_2(t,s'),
\end{aligned} \tag{A22}$$

with the initial condition

$$R_2(t,t) = L^\dagger\rho. \tag{A23}$$

The solutions for R_1, R_2 should take the following forms

$$R_1 = c_1(t,s)\sum_i a_i\rho + c_2(t,s)\sum_i \rho a_i, \tag{A24}$$

$$R_2 = c_3(t,s)\sum_i a_i^\dagger\rho + c_4(t,s)\sum_i \rho a_i^\dagger. \tag{A25}$$

Substituting Eq. (A24-A25) into Eq. (A19-A22), the equations for the coefficients can be derived as

$$\begin{aligned}\frac{\partial}{\partial s}c_1 &= -i\omega c_1 - 2i\lambda c_1 - N\int_0^s ds'\alpha_1(s,s')c_1(t,s')\\ &\quad + N\int_0^t ds'\alpha_2^*(s,s')c_4(t,s')\\ &\quad - N\int_s^t ds'\alpha_2(s',s)c_1(t,s'), \qquad \text{(A26)}\end{aligned}$$

$$\begin{aligned}\frac{\partial}{\partial s}c_2 &= -i\omega c_2 - 2i\lambda c_2 - N\int_0^s ds'\alpha_1(s,s')c_2(t,s')\\ &\quad + N\int_0^t ds'\alpha_2^*(s,s')c_3(t,s')\\ &\quad - N\int_s^t ds'\alpha_2(s',s)c_2(t,s'), \qquad \text{(A27)}\end{aligned}$$

$$\begin{aligned}\frac{\partial}{\partial s}c_3 &= i\omega c_3 + 2i\lambda c_3 + N\int_0^s ds'\alpha_2(s,s')c_3(t,s')\\ &\quad - N\int_0^t ds'\alpha_1^*(s,s')c_2(t,s')\\ &\quad + N\int_s^t ds'\alpha_1(s',s)c_3(t,s'), \qquad \text{(A28)}\end{aligned}$$

$$\begin{aligned}\frac{\partial}{\partial s}c_4 &= i\omega c_4 + 2i\lambda c_4 + N\int_0^s ds'\alpha_2(s,s')c_4(t,s')\\ &\quad - N\int_0^t ds'\alpha_1^*(s,s')c_1(t,s')\\ &\quad + N\int_s^t ds'\alpha_1(s',s)c_4(t,s'), \qquad \text{(A29)}\end{aligned}$$

with the initial conditions

$$\begin{aligned}c_1(t,t) &= 1,\\ c_2(t,t) &= 0,\\ c_3(t,t) &= 1,\\ c_4(t,t) &= 0. \qquad \text{(A30)}\end{aligned}$$

Then, substituting Eq. (A24-A25) into Eq. (A16), the final master equation will be derived as it is shown in Eq. (37). Note that the coefficients $C_i(t)$ are determined by $c_i(t,s)$ ($i = 1,2,3,4$) and the correlation functions $\alpha_j(t,s)$ ($j = 1,2$) as shown in Eq. (38).

---

[1] C. W. Gardiner and P. Zoller, *Quantum Noise* (Springer-Verlag, Berlin, 2004).

[2] H. P. Breuer and F. Petruccione, *Theory of Open Quantum Systems* (Oxford, New York, 2002).

[3] J. P. Paz and W. H. Zurek, in *Coherent Atomic Matter Waves*, edited by R. Kaiser, C. Westbrook, and F. David, Les Houches Summer School (Springer, Berlin, 2001), Session LXXII; july 27 - august 27, 1999, pp. 533–614, quant-ph/0010011.

[4] J. P. Paz and A. J. Roncaglia, Phys. Rev. Lett. **100**, 220401 (2008).

[5] T. Yu and J. H. Eberly, Phys. Rev. Lett. **93**, 140404 (2004); Phys. Rev. Lett. **97**, 140403 (2006).

[6] J. J. Halliwell and T. Yu, Phys. Rev. D **53**, 2012 (1996).

[7] J. An and W. M. Zhang, Phys. Rev. A **76**, 042127 (2007).

[8] C. H. Fleming and B. L. Hu, Ann. Phys. **327**, 1238 (2012).

[9] W. M. Zhang, M. H. Wu, C. U. Lei, and H. N. Xiong, Opt. Express **18**, 18407 (2010).

[10] H. N. Xiong, W. M. Zhang, X. G. Wang, and M. H. Wu, Phys. Rev. A **82**, 012105 (2010).

[11] K.-L. Liu and H.-S. Goan, Phys. Rev. A **76**, 022312 (2007).

[12] C. H. Chou, T. Yu and B. L. Hu, Phys. Rev. E **77**, 011112 (2008).

[13] H. Carmichael, *An Open System Approach to Quantum Optics* (Springer-Verlag, Berlin, 1994).

[14] J. Dalibard, Y. Castin, and K. Mölmer, Phys. Rev. Lett. **68**, 580 (1992).

[15] N. Gisin and I. C. Percival, J. Phys. A **25**, 5677 (1992); *ibid.* **26**, 2233 (1993).

[16] For a comprehensive review, see M. B. Plenio and P. L. Knight, Rev. Mod. Phys. **70**, 101 (1998).

[17] L. Diósi and W. T. Strunz, Phys. Lett. A **235**, 569 (1997).

[18] L. Diósi, N. Gisin, and W. T. Strunz, Phys. Rev. A **58**, 1699 (1998); W. Strunz, L. Diósi, and N. Gisin, Phys. Rev. Lett. **82**, 1801 (1999).

[19] T. Yu, L. Diósi, N. Gisin, and W. T. Strunz, Phys. Rev. A **60**, 91 (1999).

[20] X. Zhao, J. Jing, B. Corn, and T. Yu, Phys. Rev. A **84**, 032101 (2011).

[21] J. Jing *et al*, arXiv:1012.0364v1.

[22] J. Jing and T. Yu, Phys. Rev. Lett. **105**, 240403 (2010).

[23] J. Jing, X. Zhao, J. Q. You, and T. Yu, Phys. Rev. A **85** 042106 (2012)

[24] W. Shi, X. Zhao, and T. Yu, arXiv:1203.2219v1; see also M. Chen, and J. Q. You, arXiv:1203.2217v1

[25] X. Zhao, W. Shi, L.-A. Wu, and T. Yu, arXiv:1203.2220v1

[26] W. T. Strunz and T. Yu, Phys. Rev. A **69**, 052115 (2004).

[27] H. Yang, H. Miao, and Y. Chen, arXiv:1108.0963v3 (2011).

[28] J. Jing, L.-A. Wu, J. Q. You, and T. Yu, Phys. Rev. A **85**, 032123 (2012).

[29] G. Ritschel, J. Roden, W. T. Strunz, A. Aspuru-Guzik, and A. Eisfeld, arXiv:1108.3452v2.

[30] S. L. Braunstein and P. van Loock, Rev. Mod. Phys. **77**, 513 (2005).

[31] T. Pellizzari, Phys. Rev. Lett. **79**, 5242 (1997).
[32] T. Yu, Phys. Rev. A **69**, 062107 (2004).
[33] M. O. Scully and M. S. Zubairy, *Quantum Optics* (Cambridge University Press, Cambridge, England, 1997).